

Review Article



Good News about Application of Advanced Methods in Food Examination

Fahim Aziz Eldin Shaltout

Food Control Department, Faculty of Veterinary Medicine, Benha University, Egypt.

Corresponding Author: Fahim Aziz Eldin Shaltout Food Control Department, Faculty of Veterinary Medicine, Benha University, Egypt.**Received date:** November 15, 2024; **Accepted date:** November 25, 2024; **Published date:** December 15, 2024**Citation:** Fahim Aziz Eldin Shaltout, Good News about Application of Advanced Methods in Food Examination, Dietary Nourishment and Food Processing Techniques, vol 1(3). DOI: 10.9567/3064-7061/WSJ.110**Copyright:** © 2024, Fahim Aziz Eldin Shaltout, this is an open-access article distributed under the terms of The Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The Electronic noses are devices able to characterize and differentiate the aroma profiles of the various foods, especially the meat and the meat products. During the recent years advanced method, the e-noses have been widely used in the food analysis and proved to provide a fast, simple, non-expensive and non-destructive method of the food assessment and quality control. The aim of this study is to summarize the most important features of this analytic tool and to present basic fields and the typical areas of the e-nose use as well as most commonly used sensor types and patterns for the e-nose design. Prospects for the future development of this technique are presented. The Methods and the researches may be a guideline for the practical e-nose use.

Keywords: electronic nose; sensors; gas chromatography; odor.

Introduction

The most important factors affecting freshness and quality of the poultry and the fish are the color, the texture and the flavor. The Meat flavor quality is determined by composition of VOC as its critical factor. They should be qualified by electronic nose (e-nose) in the same way as human senses, because these modalities are the basis for human perception of freshness and quality. The odor of the meat is formed by a complex mixture of the different volatile organic compounds, originating from various reactions. It is often stated that the fresh meat is almost unscented and the only odor that can be detected by the consumer is described as bloody. During storage or thermal processes, the precursors of the aroma constituents undergo oxidation and decomposition reactions leading to a number of products which can then react further providing the organic compounds of low molecular mass and usually these secondary products are responsible for the odor development (1,2,3,4,5,6 and 7). The characteristic aroma of the meat usually originates from the thermal processes like the cooking, the roasting or the frying. The Chemical reactions occurring during the heat-treatment involve degradation of the amino acids, the peptides, the sugars, the ribonucleotides, the lipids and the vitamins. The key process in the aroma formation is Maillard reaction that occurs between reducing sugars and amino acids derivatives. The characteristic compounds for the cooked meat are the aldehydes, the ketones, the alcohols, the hydrocarbons, the

pyrazine derivatives and the Sulphur compounds. A specific effect is the warmed-over flavor (WOF), which develops during the storage in the meat after the cooking. It comes from the lipid oxidation induced by the iron released from the myoglobin and hemoglobin during the cooking. The unpleasant odor is described as the metallic, the musty and the pungent and the secondary oxidation products responsible for it are hexanal, 2,3-octanediol, pentanal, 2-pentylfuran and 2-octenal (8,9,10,11,12,13 and 14). Another process connected to the odor development apart from the heat treatment of the meat is spoilage. During the storage the bacterial activity leads to the production of the volatile organic compounds of the unpleasant odor, most of which are easily identifiable (methanol, ethanol, dimethyl sulfide, methyl thioacetate, toluene, nonane, 2,3-butanediol and others). The Alcohols, the hydrocarbons, the aliphatic the ketones, the volatile acids and the benzenic compounds are the compounds typical for the physicochemical changes generated as a result of fat decomposition and lipid oxidation. The aroma profile of the meat can be influenced by different factors like the fat content, the animal diet, the breed, the pH, the storage condition or the heat treatment method. This subject is still not fully explored despite of many discussions. Assessment of the meat freshness can be done by analysis of mixture of volatile organic compounds. Classical methods of analysis of the meat aroma profile involve the gas chromatography (GC) and the olfactometry (O). A fusion of these two techniques, GC-O, combines the ability of the chromatographic unit to separate different

compounds and the capability of the olfactometric system to characterize them (24,25,26,27,28,29,30,31 and 32). Apart from the chemical methods, the sensory descriptive analysis is still of great significance. The aroma profile is difficult to analyse and characterize in the laboratory since the human sense of smell does not distinguish individual components but rather identifies the specific mixture of volatiles as a whole. The Classical methods of chemical analysis based on gas chromatography result in the identification and the quantification of the particular compounds which can be considered the most important odor indicators. Often the single compounds present in meaningful quantity in the aroma profile are not noticeable by the human sense of the smell. The practice shows that there is not always a simple correlation between the concentration of particular compounds and the odor perception. Electronic noses are analytical instruments designed to mimic the work of the human sense of the smell. In this technique the analytic process does not concentrate on the identification and quantification of the components of the mixture of volatile compounds but rather on the quantitative description of the complete aroma profile, including the relationships between its components. In many researches the time change of the profile or the profile change in relation to standard profile is important. In such cases volatile profile is a fingerprint typical for certain test. The most important issues are presented below (33,34,35,36,37,38,39,40 and 41). The standard e-nose equipment is based on a series of gas sensors able to collect the chemical signal coming from the headspace and to transmit it to the electronic devices. Different types of sensors can be used for different applications, among them the most popular are electrochemical sensors like metal oxide semiconductors (MOS) or the conducting polymers (CP) and the piezoelectric sensors like quartz microbalances (QCM). The Optical, calorimetric and biosensors can be also incorporated into the e-nose system. Those innovative types of advanced method e-noses based on gas chromatography (GC) or gas chromatography with mass spectrometry (GC/MS) have been introduced recently and their high utility has been demonstrated. In this type of detection each peak from a mass spectrum or a chromatogram acts as a "sensor" providing also information about the chemical structure of the corresponding compound. Usually, it is a standard to couple e-nose with multivariate software equipped with the software tools for the chemometric interpretation of the sensor signals (42,43,44,45,46,47,48,49,50 and 51). The data collected from the sensors are analyzed using various statistical tools in order to create a numerical model of the aroma profile of the sample. When having a quantitative description of the aroma profile of samples one can then compare the unknown samples with the reference materials or study the influence of different factors on the odor. The most effective statistical methods used in these types of investigation include principal component analysis (PCA), partial least

squares regression (PLSR), linear and canonical discriminant analysis (LDA and CDA) or the artificial neural networks (ANN). Although advanced method the e-noses are rather an unreliable tool for the determination of specific chemical compounds constituting the aroma profile, the advantages of this method over the traditional ones are conspicuous. The analyses are fast, simple and low-cost, what makes the e-nose a gratifying analytical method for quality-control applications. However, this technique requires specific and time-consuming training for staff. This review paper aims to present diversity of types of advanced method the e-noses used for the evaluation of the meat and the meat products. Prospects for the future development of this technique are presented. The Methods and the researches which discussed may be a guideline for the practical e-nose application (62,63,64,65,66,67,68,69 and 70).

The Applications of the e-nose in the meat analysis

The Spoilage monitoring Traditional methods of the meat spoilage

The monitoring is based on measurements of chemical or biological spoilage indicators. Standard tests include total bacterial count (TBC), rancidity measurements with thiobarbituric acid reactive substances assay (TBARS), determination of the VOCs by GC-MS and total volatile basic nitrogen (TVBN) measurements. The Colour evaluation and sensory descriptive analysis can be also useful in the detection of spoiled samples. However, all these methods are rather costly, time consuming and require specialized staff, what makes them difficult for on-line applications (71,72,73,74,75,76,77,78,79 and 80). As the spoilage process is strongly connected with odor changes, the most of studies investigate the possibilities of employing e-nose to this purpose. A standard procedure in this type of study involves choosing the storage conditions (the type of the packaging and the temperature), usually mimicking storage in typical household or industrial conditions and analyzing samples after different periods of time. A special attention must be paid to ensure careful preparation of the samples in order to preserve the headspace composition during storage and analysis. The sensor types used for this purpose include mainly metal oxide semiconductors or conducting polymers. During last years a few applications using colorimetric sensors could be noticed (81,82,83,84,85,86,87,88 and 89). The statistical tools employed for data analysis involve multivariate statistics and artificial neuron networks. The obtained results are then confronted with data from reference methods based on the microbiological evaluation, sensory descriptive analysis or chemical determination of spoilage indicators. A number of fully successful attempts to employ advanced method e-noses for the meat spoilage monitoring prove the utility of this analytic tool. Therefore e-nose found application as a laboratory tool, yet no in industrial practice. The prospects of method development involve further investigation of the

value of information coming from particular sensors and constructing new portable instruments with reduced number of sensors in order to minimize costs and simplify the analysis (90,91,92,93,94,95,96,97,98 and 99).

The Differentiation between types of the meat

The Determination of the animal species from which the meat was produced is an important problem of the food analysis. From the consumer's point of view, it is an essential issue not only because of quality of the product or the meat adulteration but also because of health, dietary and religious aspects. The classical methods of the meat species identification involve expensive and time-consuming molecular biology-based methods or spectroscopic measurements requiring specialized equipment and complex data analysis. It has been recently demonstrated that e-nose can be a reliable analytic tool for screening of the meat samples for animal species origin (100,101,102,103,104,105,106 and 107). A successful differentiation between the meat samples from sheep, cattle, poultry, and swine was performed using an electronic nose based on gas chromatography. A simple statistical data analysis using principal component analysis (PCA) could clearly separate groups of samples derived from different animals. Moreover, although advanced method the e-noses generally do not give information about chemical composition of the headspace, the chromatographic analysis performed by this type of e-nose allowed the tentative identification of compounds responsible for the aroma profile, using Kovat's indices. The methodology presented in the study was extremely simple, fast and no-expensive, providing a useful tool for the food analysis. The ability of electronic nose to discriminate between types of the meat can be also applied in the detection of the meat adulteration with proteins from another, cheaper species. The Pork adulteration in minced mutton was studied with an e-nose containing 10 metal oxide sensors. Mutton samples were mixed with pork in different proportions and the headspace of each sample was tested with the e-nose. Various statistic tools were implemented to analyze the data obtained (PCA, DA, PLS, MLR and BPNN) and it was clearly demonstrated that the e-nose methodology coupled with multivariate analysis can easily predict the degree of adulteration of pork in minced mutton. There are also other possibilities to use e-nose technique for the meat type and quality differentiation for the meat adulteration detection and for quality control (108,109,110,111,112,113,114,115 and 116).

The Evaluation of the effect of dietary supplementation of the animals

The Animal nutrition has a potent influence on the meat quality. Different types of dietary regimes can be applied for specific purposes, among them supplementation of functional ingredients such as vitamin E, selenium, conjugated linoleic acids or omega-3 fatty acids gained a

special interest over last years. The Evaluation of the relationship between animal diet and the meat attributes is an important feature. An investigation of animal diet influence on overall antioxidant power in the meat and its connection with the aroma profile. Crossbreed steers were either pasture or grain-fed and in both groups a part of animals obtained an additional vitamin E supplementation. An e-nose with 32 conducting polymer sensors was used to analyze the aroma profile of fresh beef samples from all the four groups. Antioxidant capacity tests were also performed. The relationship between the e-nose data and the antioxidant status related variables was analyzed while the linear discriminant analysis of e-nose measurements was employed to investigate the meat samples grouping as a function of feeding (52,53,54,55,56,57,58,59,60 and 61). The e-nose distinguished correctly grain and pasture produced meat as well as supplemented and non-supplemented with vitamin E grain produced meat. These results show that the aroma profile of the meat is strongly related on the antioxidant status which affects the lipids oxidation influencing the consequent production of volatile short chained aldehydes. The e-nose proved therefore to be a useful tool to discriminate the aroma profile of the fresh meat samples with different antioxidant potential. The influence of animal dietary regimes on the lipid oxidation can manifest in the change of volatile compounds profile of the meat and this effect was also investigated with the use of e-nose. In described experiment, pigs were divided into four feeding groups with different diets: control diet, supplemental vitamin E and organic selenium diet, supplemental organic selenium diet and supplemental vitamin E diet. The samples were analyzed using electronic nose equipped with ultrafast gas chromatograph with flame-ionization detectors. The e-nose data were analyzed using the AroChemBase database and ANOVA. The measurements performed by the e-nose showed that the addition of antioxidants to the pigs feed prevented the formation of Sulphur compounds in the raw meat. The electronic nose successfully allowed to determine seventeen specific volatile compounds in the supplemented meat (15,16,17,18,19,20,21,22 and 23).

The Production process monitoring

There are rather few examples in literature of direct use of e-nose for production process monitoring. Usually these are older references such as e-nose usage to identify spoiled Iberian hams during the curing process. They discussed use of e-nose for recognition of different Iberian ham ripening times. The sensors with tin-oxide semiconductor thin films were used for the tests. Some of the sensors were doped with metal catalysts. The PCA was used for results analysis together with artificial neural network. Electronic nose can be applied to study dynamic processes occurring during production of sausages. An investigation of the seasoning processes for the dry-cured meats was performed with an electronic nose containing 12 metal oxide sensors. Fresh

pork sausages were subjected to the manufacture's protocols and 5, 7, 10, 14 days seasoning. Research included monitoring of presence of ochratoxin A-producing and non-producing *Penicillium* strains during the seasoning process. The Food contamination with the ochratoxins is very dangerous for humans and is considered as possibly carcinogenic by The International Agency for Research on Cancer in 1993. The e-nose data were analyzed using DFA. Proposed research technique was successfully applied for the rapid prediction of ochratoxin A. These mentioned applications of the e-nose are satisfactory examples of successful use of e-nose based systems for quality control (119,118,119,120,121,122,123,124 and 125).

The Specific purposes:

The boar taint and the WOF evaluation the Boar taint and the WOF are the sensory defects in the meat flavor. The Boar taint is characteristic for pork derived from non-castrated male pigs. It is connected to the presence of the androstenone and the skatole but it was shown that the sensory evaluation of the 'boar taint' level does not always agree with the absolute concentrations of these two compounds. There is therefore a demand for developing a system able to efficiently discriminate between different intensities of this undesirable odor. Used e-nose and sensory panel to measure the intensity of the boar taint in the entire male pigs. An electronic nose based on ion mobility spectrometry was employed to mimic the responses given by the sensory panel. The data from e-nose analysis were calibrated using canonical correlation with the sensory measurement and a discriminant function for separating levels of boar taint in pork by e-nose was developed (126,127,128,129,130,131 and 132). The Sensorics research affirmed stronger correlation of the boar taint with the androsterone than with the skatole. The research showed that e-nose technique based on ion mobility spectrometry may have a potential for a rapid sorting of boar fat at the slaughter line. The warmed-over flavor develops in the meat which has been pre-cooked, chill stored and reheated. The sensory analysis of volatile compounds on the meatballs derived from pigs fed with standard diet supplemented with the addition of rapeseed and palm oil using solid state-based gas sensor array system (e-nose) and the gas chromatography/gas spectrometry together with measurements of thiobarbituric acid reactive substances (TBARS). The Obtained data were analyzed using partial least square regression modelling (PLSR). MOS sensor responses showed to be significantly related to WOF characteristics detected by both sensory and chemical analysis. This shows the potential of using gas sensor technology to monitor WOF in the pork. The simple and rapid method of prediction of WOF in cooked chicken by colorimetric sensor array. The Data from colorimetric sensor array was classified using principal component analysis and hierarchical cluster analysis. Research showed that colorimetric sensor array may be successfully used to

predict WOF development in the cooked chicken meat (133,134,135,136,137 and 138)

Conclusion:

In the recent years it was demonstrated that electronic noses provide a fast, simple and non-destructive method of the meat analysis. advanced method E-noses were successfully employed mainly in the quality control of the meat being able to monitor spoilage or adulterations and the obtained results were in accordance with the sensory evaluation, offering a reliable tool for on-line analysis. The ability of e-nose to determine the antioxidant status of the meat samples was also proved. It can be stated that every factor that influences the aroma profile of the meat, could be potentially indirectly analyzed with the e-nose system and this field still seems to be insufficiently explored. Moreover, the development of new types of advanced method e-noses, based on gas chromatography, opens new perspectives for analysis of the aroma profile of the meat. Despite of many scientific works proving usefulness of e-nose based on different types of sensors or GC techniques for the meat quality appraisal or spoilage detection, use of e-nose in industry practice is insufficient. There was not found any description or report on large scale industrial application of e-nose. Almost every paper describes potentially large possibilities of use of e-nose in industrial practice, but none reports real-life implementation. This may be explained by sensor vulnerability (sensor time drift), relatively high e-nose costs and effort consuming staff training. The above does not mean that e-nose will not be used widely in future in industrial practice. There is still lack of effective application, although the potential of e-nose approach was proven.

Conflicts of Interest:

The author declare no conflicts of interest.

References:

1. Shaltout, F.A., Riad, E. M., and Abou Elhassan, Asmaa, A (2017): prevalence Of *Mycobacterium* Tuberculosis in Imported cattle offals and Its lymph Nodes. *Veterinary Medical Journal -Giza* (VMJG), 63(2): 115 – 122.
2. Cottrell T, van Peij J. Sorbitan esters and polysorbates. In: Norn V, editor. *Emulsifiers in Food Technology*. 2nd ed. Chichester: Wiley-Blackwell; 2015. pp. 271-295.
3. Shaltout, F.A., Riad, E. M., and Asmaa Abou-Elhassan (2017): Prevalence of *Mycobacterium* Spp. In Cattle Meat and Offal's Slaughtered in And Out Abattoir. *Egyptian Veterinary medical Association*, 77(2): 407 – 420.
4. Abd Elaziz, O., Fatin S. Hassanin, Fahim A. Shaltout and Othman A. Mohamed (2021):

- Prevalence of Some Foodborne Parasitic Affection in Slaughtered Animals in Local Egyptian Abattoir. *Journal of Nutrition Food Science and Technology* 2(3): 1-5.
5. Abd Elaziz, O., Fatin, S Hassanin, Fahim, A Shaltout, Othman, A Mohamed (2021): Prevalence of some zoonotic parasitic affections in sheep carcasses in a local abattoir in Cairo, Egypt. *Advances in Nutrition & Food Science* 6(2): 6(2): 25-31.
 6. Al Shorman, A. A. M.; Shaltout, F. A. and hilat, N (1999): Detection of certain hormone residues in meat marketed in Jordan. Jordan University of Science and Technology, 1st International Conference on Sheep and goat Diseases and Productivity, 23-25 October, 1999.
 7. Ebeed Saleh, Fahim Shaltout, Essam Abd Elaali (2021); Effect of some organic acids on microbial quality of dressed cattle carcasses in Damietta abattoirs, Egypt. *Damanhour Journal of Veterinary Sciences* 5(2): 17-20.
 8. Edris A, Hassanin, F. S; Shaltout, F.A., Azza H Elbaba and Nairoz M Adel (2017): Microbiological Evaluation of Some Heat-Treated Fish Products in Egyptian Markets. *EC Nutrition* 12.3 (2017): 124-132.
 9. Edris, A., Hassan, M. A., Shaltout, F.A. and Elhosseiny, S (2013): Chemical evaluation of cattle and camel meat. *Benha veterinary medical journal*, 24(2): 191-197.
 10. Edris, A. M., Hassan, M. A., Shaltout, F. A. and Elhosseiny, S (2012): Detection of E. coli and Salmonella organisms in cattle and camel meat. *Benha veterinary medical journal*, 24(2): 198-204.
 11. Edris A.M.; Hemmat M. I., Shaltout F.A.; Elshater M.A., Eman F.M.I. (2012): Study on incipient spoilage of chilled chicken cuts-up. *Benha veterinary medical journal*, 23(1), 81-86.
 12. Edris A.M.; Hemmat M.I.; Shaltout F.A.; Elshater M.A., Eman, F.M.I. (2012): Chemical analysis of chicken meat with relation to its quality. *Benha veterinary medical journal*, 23(1): 87-92 .
 13. Edris, A.M.; Shaltout, F.A. and Abd Allah, A.M. (2005): Incidence of Bacillus cereus in some meat products and the effect of cooking on its survival. *Zag. Vet. J.* 33 (2):118-124.
 14. Edris, A.M.; Shaltout, F.A. and Arab, W.S. (2005): Bacterial Evaluation of Quail Meat. *Benha Vet. Med.J.* 16 (1):1-14.
 15. Edris, A.M.; Shaltout, F.A.; Salem, G.H. and El-Toukhy, E.I. (2011): Incidence and isolation of Salmonellae from some meat products. Benha University, Faculty of Veterinary Medicine, Fourth Scientific Conference 25-27th May 2011 Veterinary Medicine and Food Safety) 172-179 benha, Egypt.
 16. Edris AA, Hassanin, F. S; Shaltout, F.A., Azza H Elbaba and Nairoz M Adel. (2017): Microbiological Evaluation of Some Heat Treated Fish Products in Egyptian Markets. *EC Nutrition* 12.3, 134-142.
 17. Edris, A.M.; Shaltout, F.A.; Salem, G.H. and El-Toukhy, E. I. (2011): Plasmid profile analysis of Salmonellae isolated from some meat products. Benha University, Faculty of Veterinary Medicine, Fourth Scientific Conference 25-27th May 2011 Veterinary Medicine and Food Safety) 194-201 benha, Egypt.
 18. Ragab A, Abobakr M. Edris, Fahim A.E. Shaltout, Amani M. Salem (2022): Effect of titanium dioxide nanoparticles and thyme essential oil on the quality of the chicken fillet. *Benha veterinary medical journal* 41(2): 38-40.
 19. Hassan, M.A, Shaltout, F. A, Arfa M. M, Mansour A.H and Saudi, K. R (2013): biochemical studies on rabbit meat related to some diseases. *Benha veterinary medical journal* 25(1):88-93.
 20. Hassan, M.A and Shaltout, F.A. (1997): Occurrence of Some Food Poisoning Microorganisms in Rabbit Carcasse, Alex. J. Vet. Science, 13(1):55-61.
 21. Hassan M, Shaltout FA and Saqur N (2020): Histamine in Some Fish Products. *Archives of Animal Husbandry & Dairy Science* 2(1): 1-3.
 22. Hassan, M.A and Shaltout, F.A. (2004): Comparative Study on Storage Stability of Beef, Chicken meat, and Fish at Chilling Temperature. *Alex. J. Vet. Science*, 20(21):21-30.
 23. Hassan, M. A; Shaltout, F. A.; Maarouf, A.A. and El-Shafey, W.S. (2014): Psychrotrophic bacteria in frozen fish with special reference to pseudomonas species. *Benha Vet. Med. J.* 27 (1):78-83.
 24. Hassan, M. A; Shaltout, F. A.; Arafa, M.M.; Mansour, A.H. and Saudi, K.R.n(2013): Bacteriological studies on rabbit meat related to some diseases. *Benha Vet. Med. J.* 25 (1):94-99.
 25. Sahi SS. (2014). Ascorbic acid and redox agents in bakery systems. In: Zhou W, editor. Bakery Products Science and Technology. 2nd ed. Chichester: Wiley-Blackwell. 183-197.
 26. Hassanin, F. S; Hassan, M. A., Shaltout, F.A., Nahla A. Shawqy and 2Ghada A. Abd-Elhameed (2017): Chemical criteria of chicken meat. *Benha veterinary medical journal*, 33(2):457-464.
 27. Hassanin, F. S; Hassan, M. A.; Shaltout, F.A. and Elrais-Amina, M (2014): clostridium perfringens in vacuum packaged meat products. *Benha veterinary medical journal*, 26(1):49-53.
 28. Hassanien, F.S.; Shaltout, F.A.; Fahmey, M.Z. and Elsukkary, H.F. (2020): Bacteriological quality guides in local and imported beef and their relation

- to public health. *Benha Veterinary Medical Journal* 39: 125-129.
29. Hassanin, F. S; Shaltout, F. A. and, Mostafa E.M(2013): Parasitic affections in edible offal. *Benha Vet. Med.J.*25 (2):34-39.
 30. Hassanin, F. S; Shaltout, F.A., Lamada, H.M., Abd Allah, E. M. (2011): the effect of preservative (nisin) on the survival of listeria monocytogenes. *Benha veterinary medical journal*, special issue [I]: 141-145.
 31. Khattab, E., Fahim Shaltout and Islam Sabik (2021): Hepatitis A virus related to foods. *Benha veterinary medical journal* 40(1): 174-179.
 32. Saad M. Saad, Fahim A. Shaltout, Amal A. A. Farag & Hashim F. Mohammed (2022): Organophosphorus Residues in Fish in Rural Areas. *Journal of Progress in Engineering and Physical Science* 1(1): 27-31.
 33. Saif, M., Saad S.M., Hassanin, F. S; Shaltout FA, Marionette Zaghloul (2019): Molecular detection of enterotoxigenic Staphylococcus aureus in ready-to-eat beef products. *Benha Veterinary Medical Journal* 37 (2019) 7-11.
 34. Saif, M., Saad S. M., Hassanin, F. S; Shaltout, F.A., Marionette Zaghlou (2019); Prevalence of methicillin-resistant Staphylococcus aureus in some ready-to-eat meat products. *Benha Veterinary Medical Journal* 37 (2019) 12-15.
 35. Farag, A. A., Saad M. Saad¹, Fahim A. Shaltout¹, Hashim F. Mohammed (2023 a): Studies on Pesticides Residues in Fish in Menofia Governorate. *Benha Journal of Applied Sciences*, 8(5): 323-330.
 36. Farag, A. A., Saad M. Saad¹, Fahim A. Shaltout¹, Hashim F. Mohammed (2023 b): Organochlorine Residues in Fish in Rural Areas. *Benha Journal of Applied Sciences*, 8 (5): 331-336.
 37. Shaltout, F.A., Mona N. Hussein, Nada Kh. Elsayed (2023): Histological Detection of Unauthorized Herbal and Animal Contents in Some Meat Products. *Journal of Advanced Veterinary Research* 13(2): 157-160.
 38. Shaltout, F. A., Heikal, G. I., Ghanem, A. M. (2022): Mycological quality of some chicken meat cuts in Gharbiya governorate with special reference to *Aspergillus flavus* virulent factors. *benha veteriv medical journal veterinary* 42(1): 12-16.
 39. Shaltout, F.A., Ramadan M. Salem, Eman M. Eldiasty, Fatma A. Diab (2022): Seasonal Impact on the Prevalence of Yeast Contamination of Chicken Meat Products and Edible Giblets. *Journal of Advanced Veterinary Research* 12(5): 641-644.
 40. Shaltout, F.A., Abdelazez Ahmed Helmy Barr and Mohamed Elsayed Abdelaziz (2022): Pathogenic Microorganisms in Meat Products. *Biomedical Journal of Scientific & Technical Research* 41(4): 32836-32843.
 41. Shaltout, F.A., Thabet, M.G. and Koura, H.A. (2017). Impact of Some Essential Oils on the Quality Aspect and Shelf Life of Meat. *J Nutr Food Sci.*, 7: 647.
 42. Shaltout, F.A., Islam Z. Mohammed², El -Sayed A. Afify (2020): Bacteriological profile of some raw chicken meat cuts in Ismailia city, Egypt. *Benha Veterinary Medical Journal* 39, 11-15.
 43. Shaltout, F.A., Islam, Z. Mohammed., El -Sayed A. Afify (2020): Detection of E. coli O157 and Salmonella species in some raw chicken meat cuts in Ismailia province, Egypt. *Benha Veterinary Medical Journal* 39 (2020) 101-104.
 44. Shaltout, F.A., E.M. El-diasty and M. A. Asmaa-Hassan (2020): hygienic quality of ready to eat cooked meat in restaurants at Cairo. *Journal of Global Biosciences* 8(12): 6627-6641.
 45. Shaltout, F.A., Marriomet Z. Nasief, L. M. Lotfy, Bossi T. Gamil (2019): Microbiological status of chicken cuts and its products. *Benha Veterinary Medical Journal* 37 (2019) 57-63.
 46. Shaltout, F.A. (2019): Poultry Meat. *Scholarly Journal of Food and Nutrition* 22 1-2.
 47. Shaltout, F.A. (2019): Food Hygiene and Control. *Food Science and Nutrition Technology* 4(5): 1-2.
 48. Hassanin, F. S; Shaltout, F.A., Seham N. Homouda and Safaa M. Arakeeb (2019): Natural preservatives in raw chicken meat. *Benha Veterinary Medical Journal* 37 (2019) 41-45.
 49. Hazaa,W., Shaltout, F.A., Mohamed El-Shate (2019): Prevalence of some chemical hazards in some meat products. *Benha Veterinary Medical Journal* 37 (2) 32-36.
 50. Hazaa,W, Shaltout, F.A., Mohamed El-Shater (2019): Identification of Some Biological Hazards in Some Meat Products. *Benha Veterinary Medical Journal* 37(2) 27-31.
 51. Gaafar, R., Hassanin, F. S; Shaltout, F.A., Marionette Zaghloul (2019): Molecular detection of enterotoxigenic Staphylococcus aureus in some ready to eat meat-based sandwiches. *Benha Veterinary Medical Journal* 37 (2) 22-26.
 52. Gaafar, R., Hassanin, F. S; Shaltout, F.A., Marionette Zaghloul (2019): Hygienic profile of some ready to eat meat product sandwiches sold in Benha city, Qalubiya Governorate, Egypt. *Benha Veterinary Medical Journal* 37 (2) 16-21.
 53. Saad S.M., Shaltout, F.A., Nahla A Abou Elroos, Saber B El-nahas (2019): Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. *J Food Sci Nutr Res.* 2 (1): 012-020.
 54. Saad S.M., Shaltout, F.A., Nahla A Abou Elroos² and Saber B El-nahas (2019): Incidence of

- Staphylococci* and *E. coli* in Meat and Some Meat Products. *EC Nutrition* 14.6 (2019).
55. Saad S.M., Hassanin, F. S.; Shaltout, F.A., Marionette Z Nassif, Marwa Z Seif. (2019): Prevalence of Methicillin-Resistant *Staphylococcus Aureus* in Some Ready-to-Eat Meat Products. *American Journal of Biomedical Science & Research* 4(6):460-464.
 56. Shaltout, Fahim (2019): Pollution of Chicken Meat and Its Products by Heavy Metals. *Research and Reviews on Healthcare: Open Access Journal*, 4, 3(381-3382).
 57. Shaltout, F. A.; E.M EL-diahy; M. S. M Mohamed (2018): Effects of chitosan on quality attributes fresh meat slices stored at 4 C. *Benha veterinary medical journal*, 35(2): 157-168.
 58. Shaltout and Abdel-Aziz, (2004): *Salmonella enterica* serovar Enteritidis in poultry meat and their epidemiology. *Vet. Med. J. Giza*, 52, pp. 429-436.
 59. Shaltout, F.A., Hala F El-Shorah, Dina I El Zahaby, Lamiaa M Lotfy (2018): Bacteriological Profile of Chicken Meat Products. *SciFed Food & Dairy Technology Journal*, 2:3.
 60. Shaltout, F.A., Mohamed, A.H. El-Shater., Wafaa Mohamed Abd El-Aziz (2015): Bacteriological assessment of Street Vended Meat Products sandwiches in kalyobia Governorate. *Benha veterinary medical journal*, 28(2), 58-66.
 61. Shaltout, F.A., Mohamed A El shatter and Heba M Fahim (2019): Studies on Antibiotic Residues in Beef and Effect of Cooking and Freezing on Antibiotic Residues Beef Samples. *Scholarly Journal of Food and Nutrition*, 2(1) 1-4.
 62. Shaltout FA, Zakaria IM and Nabil ME. (2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to *Clostridium perfringens*. *Nutrition and Food Toxicology* 2.5 (2018): 429-438.
 63. Shaltout FA, Ahmed A A Maarouf and Mahmoud ES Elkhoully. (2017): Bacteriological Evaluation of Frozen Sausage. *Nutrition and Food Toxicology* 1.5; 174-185.
 64. Shaltout FA, El-Toukhy EI and Abd El-Hai MM. (2019): Molecular Diagnosis of *Salmonellae* in Frozen Meat and Some Meat Products. *Nutrition and Food Technology Open Access* 5(1): 1-6.
 65. Shaltout, F.A., A.M.Ali and S.M.Rashad (2016): Bacterial Contamination of Fast Foods. *Benha Journal of Applied Sciences (BJAS)* 1(2)45-51.
 66. Shaltout, F.A., Zakaria. I. M., Jehan Eltanani, Asmaa. Elmelegy (2015): Microbiological status of meat and chicken received to university student hostel. *Benha veterinary medical journal*, 29(2):187-192.
 67. Saad, S. M.; Edris, A.M.; Shaltout, F. A. and Edris, Shimaa (2012): Isolation and identification of salmonellae and *E. coli* from meat and poultry cuts by using A. multiplex PCR. *Benha Vet. Med. J.* special issue 16-26.
 68. Saad, S.M. and Shaltout, F.A. (1998): Mycological Evaluation of camel carcasses at Kalyobia Abattoirs. *Vet. Med. J. Giza*, 46(3):223-229.
 69. Saad S. M., Shaltout, F.A., Nahla A Abou Elroos, Saber B El-nahas. (2019): Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. *J Food Sci Nutr Res.* 2019; 2 (1): 012-020.
 70. Saad S.M., Hassanin, F. S; Shaltout, F.A., Marionette Z Nassif, Marwa Z Seif. (2019): Prevalence of Methicillin-Resistant *Staphylococcus Aureus* in Some Ready-to-Eat Meat Products. *American Journal of Biomedical Science & Research*, 4(6):460-464.
 71. Saad S.M., Shaltout, F.A., Nahla A Abou Elroos and Saber B El-nahas. (2019): Incidence of *Staphylococci* and *E. coli* in Meat and Some Meat Products. *EC Nutrition* 14.6 (2019).
 72. Shaltout FA, Riad EM, TES Ahmed and Abou Elhassan A. (2017): Studying the Effect of Gamma Irradiation on Bovine Offal's Infected with *Mycobacterium tuberculosis* Bovine Type. *Journal of Food Biotechnology Research* 1 (6): 1-5.
 73. Shaltout FA, Zakaria IM and Nabil ME. (2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to *Clostridium perfringens*. *Nutrition and Food Toxicology* 2.5 (2018): 429-438.
 74. Shaltout FA, Mohamed, A. Hassan and Hassanin, F. S (2004): Thermal inactivation of enterohaemorrhagic *escherichia coli* o157:h7 and its sensitivity to nisin and lactic acid cultures. *1st Ann. Confr., FVM., Moshtohor, Sept, 2004*.
 75. Shaltout FA, El-diahy, E. M.; Elmesalamy, M. and Elshaer, M. (2014): Study on fungal contamination of some chicken meat products with special reference to 2 the use of PCR for its identification. Conference, Veterinary Medical Journal – Giza, vol.60: 1-10.
 76. shaltout, F.A. (2002): Microbiological Aspects of Semi-cooked chicken Meat Products. *Benha Veterinary Medical Journal* 13, 2; 15-26.
 77. Shaltout FA, Thabet, M.G2 and Hanan, A. Koura3. (2017): Impact of some essential oils on the quality aspect and shelf life of meat. *Benha veterinary medical journal*, 33, (2): 351-364.
 78. Shaltout FA, Mohammed Farouk; Hosam A.A. Ibrahim and Mostafa E.M. Afifi. (2017): Incidence of Coliform and *Staphylococcus aureus* in ready to eat fast foods. *Benha veterinary medical journal*, 32(1): 13 - 17.

79. Shaltout, F.A., Zakaria, I.M., Nabil, M.E. (2017): Detection and typing of *Clostridium perfringens* in some retail chicken meat products. *Benha veterinary medical journal*, 33(2): 283-291.
80. Shaltout, F.A. (1992): Studies on Mycotoxins in Meat and Meat by Products. M.V.Sc Thesis Faculty of Veterinary Medicine, Moshtohor, Zagazig university Benha branch.
81. Shaltout, F.A. (1996): Mycological and Mycotoxicological profile Of Some Meat products. Ph.D. Thesis, Faculty of Veterinary Medicine, Moshtohor, Zagazig University Benha branch.
82. Shaltout, F.A. (1998): Proteolytic Psychrotrophes in Some Meat products. *Alex. Vet. Med. J.* 14 (2):97-107.
83. Shaltout, F.A. (1999): Anaerobic Bacteria in Vacuum Packed Meat Products. *Benha Vet. Med.J.* 10 (1):1-10.
84. Shaltout, F.A. (2000): Protozoal Foodborne Pathogens in some Meat Products. *Assiut Vet. Med. J.* 42 (84):54-59.
85. Shaltout, F.A. (2001): Quality evaluation of sheep carcasses slaughtered at Kalyobia abattoirs. *Assiut Veterinary Medical Journal*, 46(91):150-159.
86. Shaltout, F.A. (2002): Microbiological Aspects of Semi-cooked Chicken Meat Products. *Benha Vet. Med. J.* 13(2):15-26.
87. Shaltout, F.A. (2003): *Yersinia Enterocolitica* in some meat products and fish marketed at Benha city. The Third international conference Mansoura 29-30 April.
88. Shaltout, F.A. (2009): Microbiological quality of chicken carcasses at modern Poultry plant. The 3rd Scientific Conference, Faculty of Vet. Med., Benha University, 1-3 january.
89. Shaltout, F. A. and Abdel Aziz, A.M. (2004): Salmonella enterica Sero var Enteritidis in Poultry Meat and their Epidemiology ,*Vet. Med. J., Giza*,52(3):429-436.
90. Shaltout, F. A. and Abdel Aziz, A. M. (2004): escherichia coli strains in slaughtered animals and their public health importance. *J. Egypt. Vet. Med. Association* 64(2):7-21.
91. Shaltout,F.A., Amin, R., Marionet , Z., Nassif and Shimaa, Abdel-wahab (2014): Detection of aflatoxins in some meat products. *Benha veterinary medical journal*, 27(2) :368-374.
92. Shaltout, F.A. and Afify, Jehan Riad, EM and Abo Elhasan, Asmaa, A. (2012): Improvement of microbiological status of oriental sausage. *Journal of Egyptian Veterinary Medical Association* 72(2):157-167.
93. Shaltout, F.A. and Daoud, J. R. (1996): Chemical analytical studies on rabbit meat and liver. *Benha Vet. Med.J.* 8 (2):17-27.
94. Shaltout,F.A. and Edris, A.M.(1999): Contamination of shawerma with pathogenic yeasts. *Assiut Veterinary Medical Journal*, 40(64):34-39.
95. Shaltout, F. A.; Eldiasty, E. and Mohamed, M.S. (2014): Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. Animal Health Research Institute: First International Conference on Food Safety and Technology 19-23 June 2014 Cairo Egypt pages 79-89.
96. Shaltout, F. A.; Eldiasty, E.; Salem, R. and Hassan, Asmaa (2016): Mycological quality of chicken carcasses and extending shelf – life by using preservatives at refrigerated storage. *Veterinary Medical Journal -Giza (VMJG)*62(3)1-7.
97. Shaltout, F.A.; Salem, R. Eldiasty, E.; and Diab, Fatema. (2016): Mycological evaluation of some ready to eat meat products with special reference to molecular chacterization. *Veterinary Medical Journal -Giza* 62(3)9-14.
98. Shaltout, F. A.; Elshater, M. and Wafaa, Abdelaziz (2015): Bacteriological assessment of street vended meat products sandwiches in Kalyobia Governorate. *Benha Vet. Med.J.*28 (2):58-66.
99. Shaltout, F. A.; Gerges, M.T. and Shewail, A. A. (2018): Impact of Organic Acids and Their Salts on Microbial Quality and Shelf Life of Beef. *Assiut veterinary medical journal* 64(159): 164-177.
100. Almeida EL, Chang YK. (2012). Effect of the addition of enzymes on the quality of frozen pre-baked French bread substituted with whole wheat flour. *LWT-Food Science and Technology.* 49:64-72.
101. Shaltout, F. A.; Ghoneim, A.M.; Essmail, M.E. and Yousseif, A. (2001): Studies on aflatoxin B1 residues in rabbits and their pathological effects. *J. Egypt. Vet. Med. Association* 61(2):85-103.
102. Shaltout, F. A. and Hanan, M.T. El-Lawendy (2003): Heavy Metal Residues in Shawerma. *Beni-Suef Vet. Med. J.* 13(1):213-224.
103. Shaltout, F.A. and Hashim, M.F. (2002): Histamine in salted, Smoked and Canned Fish products. *Benha Vet. Med.J.*13 (1):1-11.
104. Shaltout, F. A.; Hashim,M.F. and Elnahas, s. (2015): Levels of some heavy metals in fish (tilapia nilotica and Claris lazera) at Menufia Governorate. *Benha Vet. Med.J.*29 (1):56-64.
105. Shaltout, F.A. and Ibrahim, H.M. (1997): Quality evaluation of luncheon and Alexandrian sausage. *Benha Vet. Med.J.*10 (1):1-10.
106. Gaupp R, Adams W. (2015). Diacetyl Tartaric Acids of Monoglycerides (DATEM) and associated emulsifiers in bread making. In: Norn V, editor. Emulsifiers in Food Technology. 2nd ed. Chichester: Wiley-Blackwell, 121-145.

107. Shaltout, F.A.; Nassif, M and Shakran, A (2014): Quality of battered and breaded chicken meat products. *Global Journal of Agriculture and Food Safety Science* – 1(2) ISSN 2356-7775.
108. Shaltout, F. A., Amani M. Salem, A. H. Mahmoud, K. A (2013): Bacterial aspect of cooked meat and offal at street vendors level. *Benha veterinary medical journal*, 24(1): 320-328.
109. Shaltout, F. A. and Salem, R.M. (2000): Moulds, aflatoxin B1 and Ochratoxin A in Frozen Livers and meat products. *Vet . Med. J. Giza* 48(3):341-346.
110. Yasser H. Al-Tarazi, A. Al-Zamil, Shaltout FA. and H. Abdel- Samei (2002). Microbiological status of raw cow milk marketed in northern Jordan. *AVMJ* Volume 49 Issue 96 Pages 180-194
111. Shaltout FA, Zakaria IM and Nabil ME. (2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to *Clostridium perfringens*. *Nutrition and Food Toxicology*, 2(5):429-438.
112. Shaltout, F. A.; El-diasy, E.M. and Mohamed, M. S. (2014): Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. 1st Scientific conference of food safety and Technology, pp. 79-89.
113. Shaltout, F. A.; El-diasy, E.M.; Salem, R. M. and Asmaa, M. A. Hassan. (2016): Mycological quality of chicken carcasses and extending shelf -life by using preservatives at refrigerated storage. *Veterinary Medical Journal – Giza* ,62(3) :1-10.
114. Shaltout FA, R.M. Salem, E.M. El-Diasy and W.I.M. Hassan. (2019): Effect of Lemon Fruits and Turmeric Extracts on Fungal Pathogens in Refrigerated Chicken Fillet Meat. *Global Veterinaria* 21 (3): 156-160.
115. Shaltout FA, El-diasy, E, M.; Elmesalamy, M. and Elshaer, M. (2014): Study on fungal contamination of some chicken meat products with special reference to 2 the use of PCR for its identification. Conference, *Veterinary Medical Journal – Giza* vol. December 2014/12/17 vol.60 1-10.
116. Shaltout, F. A.; Salem, R. M; El-diasy, Eman and Fatema, A.H. Diab. (2016): Mycological evaluation of some ready to eat meat products with special reference to molecular characterization. *Veterinary Medical Journal – Giza*. 62(3): 9-14.
117. Shaltout FA, Ahmed, A.A. Maarouf, Eman, M.K. Ahmed (2018): Heavy Metal Residues in chicken cuts up and processed chicken meat products. *Benha veterinary medical journal*, 34(1): 473-483.
118. Shaltout, F.A.; Hanan M. Lamada, Ehsan A.M. Edris. (2020): Bacteriological examination of some ready to eat meat and chicken meals. *Biomed J Sci & Tech Res.*, 27(1): 20461-20465.
119. Sobhy, Asmaa and Shaltout, Fahim (2020): Prevalence of some food poisoning bacteria in semi cooked chicken meat products at Qaliubiya governorate by recent Vitek 2 compact and PCR techniques. *Benha Veterinary Medical Journal* 38 (2020) 88-92.
120. Sobhy, Asmaa and Shaltout, Fahim (2020): Detection of food poisoning bacteria in some semi-cooked chicken meat products marketed at Qaliubiya governorate. *Benha Veterinary Medical Journal*, 38 (2020) 93-96.
121. Shaltout, F.A. (2024): Abattoir and Bovine Tuberculosis as A Reemerging Foodborne Disease. *Clinical Medical Reviews and Report* 6(1):1-7.
122. Shaltout, F.A. (2023): Viruses in Beef, Mutton, Chevon, Venison, Fish and Poultry Meat Products. *Food Science & Nutrition Technology* 8(4):1-10.